


Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/Figures/Figure 15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: [REDACTED]



0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/Figure 15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: :



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210896_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: [REDACTED]



0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: [REDACTED]



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/acripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number:



0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: [REDACTED]



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number: 7



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TEL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number:



⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TPL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number:



⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20

Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 517 of 658

Merck Serono
EMR 200125-002

Page 35 of 42
Confidential

Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values)
(Safety Population)

Subject ID/Random Number: 1



⊖: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Merck Serono
EMR 200125-002
Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



8: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/Figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.9 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



⊖: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.9.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
⊖: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: i



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/Figure 15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 538 of 658


Merck Serono
EMR 200125-002

Page 14 of 42

Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: .



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL_programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 539 of 658

Merck Serono
EMR 200125-002

Page 15 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20




Levothyroxine
EMR 200125-002

Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 540 of 658

Page 16 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: [REDACTED]



Treatment: ●—● Treatment A

0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/Figures/Figure 15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002


Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 542 of 658

Page 18 of 42
Confidential

Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—●—● Treatment A ◆—◆—◆ Treatment B ▲—▲—▲ Treatment C
0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 543 of 658

Merck Serono
EMR 200125-002

Page 19 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1




Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: 1



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 546 of 658


Page 22 of 42

Merck Serono
EMR 200125-002

Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ◆◆◆ Treatment B


@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TEL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 549 of 658

Merck Serono
EMR 200125-002

Page 25 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

@:Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


EMR200125-002 Page 550 of 658
ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted

Merck Serono
EMR 200125-002

Page 26 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/Figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 553 of 658

Merck Serono
EMR 200125-002

Page 29 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

⊖: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:




Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—●—● Treatment A ◆—◆—◆ Treatment B ▲—▲—▲ Treatment C

⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure 15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ◆◆◆ Treatment B

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL_programs/Figures/Figure 15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 560 of 658

Merck Serono
EMR 200125-002

Page 36 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(For Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

8: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (For Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●●● Treatment A ◆◆◆ Treatment B ▲▲▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportion:
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 563 of 658

Merck Serono
EMR 200125-002

Page 39 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

⊘: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions
State


EMR200125-002 Page 564 of 658
ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted

Merck Serono
EMR 200125-002

Page 40 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale
(for Total T4 Values) (Safety Population)

Subject ID/Random Number: .



Treatment: ▲▲▲ Treatment C

0: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

Source: listing 16.2.5.2


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/Figures/Figure
15.4.2.10.sas

10DEC2014 00:20



Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number: .



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C
@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002


Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 566 of 658

Merck Serono
EMR 200125-002

Page 42 of 42
Confidential

Figure 15.4.2.10 Concentration-Time Profiles: Individual Concentration versus Actual Time in Semi-Logarithmic Scale (for Total T4 Values) (Safety Population)

Subject ID/Random Number:



Treatment: ●—● Treatment A ◆—◆ Treatment B ▲—▲ Treatment C

@: Subjects excluded from PK Population; Horizontal lines represent the Baseline of the respective treatments.

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


Source: listing 16.2.5.2

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.10.sas

10DEC2014 00:20

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: .



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TEL programs/figures/figure 15.4.2.11.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210896_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: ...




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values)
(Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population:
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002


Dosage Form Proportion: ... ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 573 of 658

Merck Serono
EMR 200125-002

Page 7 of 42
Confidential

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values)
(Safety Population)

Subject ID/Random Number: :



0: Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 11




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 1



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure 15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas


10DEC2014 00:20



Merck Serono
EMR 200125-002

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure 15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 11



@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions
State


EMR200125-002 Page 580 of 658 (50 µg, 100 µg, and 200 µg Tablets) in Fasted State

Merck Serono
EMR 200125-002

Page 14 of 42
Confidential

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 1




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 1



@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions: 50 µg, 100 µg, and 200 µg Tablets in Fasted State
EMR200125-002 Page 582 of 658


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State

Merck Serono
EMR 200125-002

Page 16 of 42
Confidential

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: 1



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure 15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@: Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Merck Serono
EMR 200125-002
Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (For Total T3 Values)
(Safety Population)

Subject ID/Random Number:




@:Subjects excluded from FK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values)
(Safety Population)

Subject ID/Random Number: 1




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TPL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20

Levothyroxine
EMR 200125-002

Dosage Form Proportions
State


ions (50 µg, 100 µg, and 200 µg Tablets) in Fasted
EMR200125-002 Page 589 of 658

Merck Serono
EMR 200125-002

Page 23 of 42
Confidential

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values)
(Safety Population)

Subject ID/Random Number:



@: Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




0: Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;

Treatment C: 3 tablets of 200 µg of levothyroxine.


/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values)
(Safety Population)

Subject ID/Random Number:




@: Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/Figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: .



@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Levothyroxine
EMR 200125-002

Dosage Form Proportions (50 µg, 100 µg, and 200 µg Tablets) in Fasted State
EMR200125-002 Page 594 of 658


Page 28 of 42

Merck Serono
EMR 200125-002

Confidential

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population;

Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;


Treatment C: 3 tablets of 200 µg of levothyroxine.

/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure 15.4.2.11.sas

10DEC2014 00:20

Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (For Total T3 Values)
(Safety Population)

Subject ID/Random Number: :




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number: :




@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20



Figure 15.4.2.11 Concentration-Time Profiles: Individual Concentration versus Actual Time in Linear Scale (for Total T3 Values) (Safety Population)

Subject ID/Random Number:



@:Subjects excluded from PK Population;
Treatment A: 12 tablets of 50 µg of levothyroxine; Treatment B: 6 tablets of 100 µg of levothyroxine;
Treatment C: 3 tablets of 200 µg of levothyroxine.
/project24/ep/blinded/e210898_merc/stats/versioncontrol/primary/scripts/program/main/TFL programs/figures/figure
15.4.2.11.sas

10DEC2014 00:20

