

**Fig. 30a: Concentration of CPA in the serum of subject 30 following oral administration of one tablet of Androcur 100 mg.**  
Top: linear presentation; bottom: semilogarithmic presentation




Fig. 31a: Concentration of CPA in the serum of subject 31 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation





Fig. 32a: Concentration of CPA in the serum of subject 32 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation



**Fig. 33a:** Concentration of CPA in the serum of subject 33 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation




Fig. 34a: Concentration of CPA in the serum of subject 34 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation




Fig. 35a: Concentration of CPA in the serum of subject 35 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation




Fig. 36a: Concentration of CPA in the serum of subject 36 following oral administration of one tablet of Androcur 100 mg.  
Top: linear presentation; bottom: semilogarithmic presentation




Fig. 1b: Concentration of CPA in the serum of subject 1 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 2b: Concentration of CPA in the serum of subject 2 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 3b: Concentration of CPA in the serum of subject 3 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 4b: Concentration of CPA in the serum of subject 4 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation

**Fig. 5b is not available: subject 5 was dropped from the trial 7 hours after administration of two tablets Androcur 50.**





Fig. 6b: Concentration of CPA in the serum of subject 6 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 7b: Concentration of CPA in the serum of subject 7 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 8b: Concentration of CPA in the serum of subject 8 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 9b: Concentration of CPA in the serum of subject 9 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation



**Fig. 10b: Concentration of CPA in the serum of subject 10 following oral administration of two tablets of Androcur 50 mg.**  
Top: linear; bottom: semilogarithmic presentation




Fig. 11b: Concentration of CPA in the serum of subject 11 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 12b: Concentration of CPA in the serum of subject 12 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 13b: Concentration of CPA in the serum of subject 13 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 14b: Concentration of CPA in the serum of subject 14 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 15b: Concentration of CPA in the serum of subject 15 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 16b: Concentration of CPA in the serum of subject 16 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 17b: Concentration of CPA in the serum of subject 17 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 18b: Concentration of CPA in the serum of subject 18 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 19b: Concentration of CPA in the serum of subject 19 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 20b: Concentration of CPA in the serum of subject 20 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation



**Fig. 21b: Concentration of CPA in the serum of subject 21 following oral administration of two tablets of Androcur 50 mg.**  
Top: linear; bottom: semilogarithmic presentation




Fig. 22b: Concentration of CPA in the serum of subject 22 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 23b: Concentration of CPA in the serum of subject 23 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 24b: Concentration of CPA in the serum of subject 24 following oral administration of two tablets of Androcur 50 mg.

Top: linear; bottom: semilogarithmic presentation




Fig. 25b: Concentration of CPA in the serum of subject 25 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 26b: Concentration of CPA in the serum of subject 26 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 27b: Concentration of CPA in the serum of subject 27 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 28b: Concentration of CPA in the serum of subject 28 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation



**Fig. 29b: Concentration of CPA in the serum of subject 29 following oral administration of two tablets of Androcur 50 mg.**  
Top: linear; bottom: semilogarithmic presentation




Fig. 30b: Concentration of CPA in the serum of subject 30 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 31b: Concentration of CPA in the serum of subject 31 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 32b: Concentration of CPA in the serum of subject 32 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 33b: Concentration of CPA in the serum of subject 33 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig.34b: Concentration of CPA in the serum of subject 34 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation





Fig. 35b: Concentration of CPA in the serum of subject 35 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation




Fig. 36b: Concentration of CPA in the serum of subject 36 following oral administration of two tablets of Androcur 50 mg.  
Top: linear; bottom: semilogarithmic presentation