

2.7.5 References

1. WHO monographs on medicinal plants commonly used in the Newly Independent States (NIS). [Geneva, Switzerland]: World Health Organization; 2010.
2. E/S/C/O/P monographs : the scientific foundation for herbal medicinal products. 2nd ed., completed rev. and expanded. ed. Exeter: ESCOP; 2003.
3. Kitajima J, Ishikawa T, Urabe A, Satoh M. Monoterpenoids and their glycosides from the leaf of thyme. *Phytochemistry*. 2004;65(24):3279-87.
4. Takeuchi H, Lu ZG, Fujita T. New monoterpene glucoside from the aerial parts of thyme (*Thymus vulgaris* L.). *Biosci Biotechnol Biochem*. 2004;68(5):1131-4.
5. European Medicines Agency. Assessment report on *Thymus vulgaris* L., *vulgaris zygis* L., herba (EMA/HMPC/342334/2013). 2013.
6. Kohlert C, Schindler G, Marz RW, Abel G, Brinkhaus B, Derendorf H, et al. Systemic availability and pharmacokinetics of thymol in humans. *J Clin Pharmacol*. 2002;42(7):731-7.
7. Barnes J, Anderson LA, Phillipson JD. Herbal medicines. 3rd ed. London ; Grayslake, IL: Pharmaceutical Press; 2007. x, 710 p. p.
8. Knols G, Stal PC, van Ree JW. Produktieve hoest: tijm of broomhexine? Een dubbelblind gerandomiseerd onderzoek. *Huisarts en Wetenschap*. 1994;37:392-4.
9. Basch E, Ulbricht C, Hamneress P, Bevins A, Sollars D. Thyme (*Thymus vulgaris* L.), Thymol. *J Herb Pharmacother*. 2004;4(1):49-67.
10. Ernst E, Marz R, Sieder C. A controlled multi-centre study of herbal versus synthetic secretolytic drugs for acute bronchitis. *Phytomedicine*. 1997;4(4):287-93.
11. Kemmerich B, Eberhardt R, Stammer H. Efficacy and tolerability of a fluid extract combination of thyme herb and ivy leaves and matched placebo in adults suffering from acute bronchitis with productive cough. A prospective, double-blind, placebo-controlled clinical trial. *Arzneimittelforschung*. 2006;56(9):652-60.
12. Kemmerich B. Evaluation of efficacy and tolerability of a fixed combination of dry extracts of thyme herb and primrose root in adults suffering from acute bronchitis with productive cough. A prospective, double-blind, placebo-controlled multicentre clinical trial. *Arzneimittelforschung*. 2007;57(9):607-15.
13. Gruenwald J, Graubaum HJ, Busch R. Efficacy and tolerability of a fixed combination of thyme and primrose root in patients with acute bronchitis. A double-blind, randomized, placebo-controlled clinical trial. *Arzneimittelforschung*. 2005;55(11):669-76.
14. Büechi S, Vogelin R, von Eiff MM, Ramos M, Melzer J. Open trial to assess aspects of safety and efficacy of a combined herbal cough syrup with ivy and thyme. *Forsch Komplementarmed Klass Naturheilkd*. 2005;12(6):328-32.
15. Foster BC, Vandenhoek S, Hana J, Krantis A, Akhtar MH, Bryan M, et al. In vitro inhibition of human cytochrome P450-mediated metabolism of marker substrates by natural products. *Phytomedicine*. 2003;10(4):334-42.
16. Sasaki K, Wada K, Tanaka Y, Yoshimura T, Matuoka K, Anno T. Thyme (*Thymus vulgaris* L.) leaves and its constituents increase the activities of xenobiotic-metabolizing enzymes in mouse liver. *J Med Food*. 2005;8(2):184-9.

17. Aydin S, Basaran AA, Basaran N. The effects of thyme volatiles on the induction of DNA damage by the heterocyclic amine IQ and mitomycin C. Mutat Res. 2005;581(1-2):43-53.

Mechelen, 2014-12-03

Name of expert: Robert Stark